

Symmetry in the Coplanarity Condition

We can rewrite the triple product without difficulty using

$$t = \mathring{r}\mathring{d} \cdot \mathring{q}\mathring{\ell} = \mathring{r} \cdot \mathring{q}\mathring{\ell}\mathring{d}^* = \mathring{q}^*\mathring{r} \cdot \mathring{\ell}\mathring{d}^*. \quad (1)$$

Noting that $\mathring{\ell}^* = -\mathring{\ell}$ and $\mathring{r}^* = -\mathring{r}$, since \mathring{r} and $\mathring{\ell}$ are quaternions with zero scalar parts, we first obtain

$$\boxed{t = \mathring{r}\mathring{q} \cdot \mathring{d}\mathring{\ell}} \quad (2)$$

We then find by expanding the dot-product for t in terms of the scalar and vector components of $\mathring{q} = (q, \mathbf{q})$ and $\mathring{d} = (d, \mathbf{d})$:

$$(\mathbf{d} \cdot \mathbf{r})(\mathbf{q} \cdot \mathbf{\ell}) + (\mathbf{q} \cdot \mathbf{r})(\mathbf{d} \cdot \mathbf{\ell}) + (dq - \mathbf{d} \cdot \mathbf{q})(\mathbf{\ell} \cdot \mathbf{r}) + d[\mathbf{r} \mathbf{q} \mathbf{\ell}] + q[\mathbf{r} \mathbf{d} \mathbf{\ell}]. \quad (3)$$

While

$$\mathring{s} = \sum_{i=1}^n w_i e_i (\mathring{r}_i \mathring{d}_i \mathring{\ell}_i^*) \quad \text{and} \quad \mathring{t} = \sum_{i=1}^n w_i e_i (\mathring{r}_i^* \mathring{q}_i \mathring{\ell}_i). \quad (4)$$

We also still have the three equations

$$\mathring{q} \cdot \delta \mathring{q} = 0, \quad \mathring{d} \cdot \delta \mathring{d} = 0, \quad \text{and} \quad \mathring{q} \cdot \delta \mathring{d} + \mathring{d} \cdot \delta \mathring{q} = 0, \quad (5)$$

all of which we can shuffled around into matrix form

$$\begin{pmatrix} A & B & \mathring{q} & 0 & \mathring{d} \\ B^T & C & 0 & \mathring{d} & \mathring{q} \\ \mathring{q}^T & 0^T & 0 & 0 & 0 \\ 0^T & \mathring{d}^T & 0 & 0 & 0 \\ \mathring{d}^T & \mathring{q}^T & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \delta \mathring{d} \\ \delta \mathring{q} \\ \lambda \\ \mu \\ \nu \end{pmatrix} = - \begin{pmatrix} \mathring{s} \\ \mathring{t} \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad (6)$$

Note that the upper left 8×8 sub-matrix is the *weighted* sum of flattened dyadic products (as first shown by Žari, Bārūk, and Łolaž)

$$\sum_{i=1}^n w_i \vec{c}_i \vec{c}_i^T, \quad (7)$$

where the eight component vector \vec{c}_i is given by

$$\vec{c}_i = \begin{pmatrix} \mathring{r}_i \mathring{d}_i \mathring{\ell}_i^* \\ \mathring{r}_i^* \mathring{q}_i \mathring{\ell}_i \end{pmatrix} = - \begin{pmatrix} \mathring{r}_i \mathring{q}_i \mathring{\ell}_i \\ \mathring{r}_i \mathring{d}_i \mathring{\ell}_i \end{pmatrix}. \quad (8)$$

We conclude that the number of solutions is equal to the number of ways of partitioning the set of variables, namely

$$\binom{n+m-2}{n-1} = \binom{n+m-2}{m-1} = \frac{(n+m-2)!}{(n-1)!(m-1)!} \quad (9)$$

To implement the numerical solution, take a small step $\delta\lambda$ in λ and solve for the increment $\delta\mathbf{x}$ in

$$\frac{d\mathbf{h}}{d\lambda} \delta\lambda + \frac{d\mathbf{h}}{d\mathbf{x}} \delta\mathbf{x} = 0, \quad (10)$$

where $J = (d\mathbf{h}/d\mathbf{x})$ is the Jacobian of \mathbf{h} with respect to \mathbf{x} .